

Estudios de Biofeedback relacionados con artritis

Astin, J.A., Beckner, W., Soeken, K., Hochberg, M.C., & Berman, B. (2002). Psychological interventions for rheumatoid arthritis: A meta-analysis of randomized controlled trials. *Arthritis and Rheumatism*, 47(3), 291-302

Bradley, L.A. (1985). Effects of cognitive-behavioral therapy on pain behavior of rheumatoid arthritis (RA) patients: Preliminary outcomes. *Scandinavian Journal of Behaviour Therapy*, 14(2), 51-64.

Bradley, L.A., Young, L.D., Anderson, K.O., Turner, R.A., Agudelo, C.A., McDaniel, L.K., et al. (1987). Effects of psychological therapy on pain behavior of rheumatoid arthritis patients: Treatment outcome and six-month follow up. *Arthritis and Rheumatism*, 30(10), 1105-1114.

Flor, H., Haag, G., & Turk, D.C. (1986). Long-term efficacy of EMG biofeedback for chronic rheumatic back pain. *Pain*, 27(2), 195-202.

Flor, H., Haag, G., Turk, D.C., & Koehler, H. (1983). Efficacy of EMG biofeedback, pseudotherapy, and conventional medical treatment for chronic rheumatic back pain. *Pain*, 17(1), 21-31.

Lavigne, J.V., Ross, C.K., Berry, S.L., & Hayford, J.R. (1992). Evaluation of a psychological treatment package for treating pain in juvenile rheumatoid arthritis. *Arthritis Care & Research*, 5(2), 101-110.

Young, L.D., Bradley, L.A., & Turner, R.A. (1995). Decreases in health care resource utilization in patients with rheumatoid arthritis following a cognitive behavioral intervention. *Biofeedback and Self-Regulation*, 20(3), 259-268.